

Flux fluctuations in a multi-random-walker model and surface growth dynamics

S. Y. Yoon, Byoung-sun Ahn and Yup Kim*

Department of Physics and Research Institute for Basic Sciences, Kyung Hee University, Seoul 130-701, Korea

We study the dynamics of visitation flux in a multi-random-walker model by comparison to surface growth dynamics in which one random walker drops a particle to a node at each time the walker visits the node. In each independent experiment (trial or day) for the multi-random-walker model, the number of walkers are randomly chosen from the uniform distribution $[\langle N_{RW} \rangle - \Delta N_{RW}, \langle N_{RW} \rangle + \Delta N_{RW}]$. The averaged fluctuation $\bar{\sigma}(T_{RW})$ of the visitations over all nodes i and independent experiments is shown to satisfy the power-law dependence on the walk step T_{RW} as $\bar{\sigma}(T_{RW}) \simeq T_{RW}^\beta$. Furthermore two distinct values of the exponent β are found on a scale-free network, a random network and regular lattices. One is β_i , which is equal to the growth exponent β for the surface fluctuation W in one-random-walker model, and the other is $\beta = 1$. β_i is found for small ΔN_{RW} or for the system governed by the internal intrinsic dynamics. In contrast $\beta = 1$ is found for large ΔN_{RW} or for the system governed by the external flux variations. The implications of our results to the recent studies on fluctuation dynamics of the nodes on networks are discussed.

PACS numbers: 05.40.-a, 89.75.-k, 89.75.Da

Recent advances of network theories [1] have elucidated that the dynamical structure of many complex systems from the internet through the biosystems to the social systems forms nontrivial networks such as scale-free networks. Therefore the dynamical behaviors of the interacting (linked) units (nodes) on the networks are very important to understand the dynamics of such complex systems. Recently an important theory [2, 3, 4, 5, 6] on the dynamical behaviors on such interacting (correlated) nodes is suggested and confirmed on the internet, World Wide Web(WWW), river networks and etc.

The main theory [2, 3] for the dynamics of the node is that there exists the power-law relation between the average activity or flux $\langle f_i \rangle$ of each node i to the flux fluctuation σ_i as

$$\sigma_i \sim \langle f_i \rangle^\alpha. \quad (1)$$

Furthermore the theory suggested that there can be two distinct classes of dynamical systems. One class consists of the systems with $\alpha \simeq 1/2$ and the other consists of those with $\alpha = 1$. The $\alpha \simeq 1/2$ systems are those in which the main controlling dynamics is the internal intrinsic dynamics. Examples are signal activity in microprocessors and time-resolved information in internet routers. The $\alpha = 1$ systems are those in which the main dynamics is controlled by the external variations. Examples are river networks, highway traffic and World Wide Webs. Recently another study [4] suggested a dynamical model which has nonuniversal α values on complex networks by introducing an impact variable to each node.

To understand the origin of two distinct classes, several dynamical models have been suggested [2, 3]. One of them is based on a multi-random-walker model. The details of the multi-random-walker model are as follows. In each experiment or trial d , which was called as "day"

in Refs. [2, 3], the number of unbiased random walkers N_{RW} are randomly chosen from the uniform distribution $[\langle N_{RW} \rangle - \Delta N_{RW}, \langle N_{RW} \rangle + \Delta N_{RW}]$. The walkers are initially randomly distributed on a given support (network) with the N nodes (sites). Then the distributed walkers simultaneously take preassigned fixed T_{RW} steps in a given experiment d and count the number of visits (or flux) f_{id} to the node i during T_{RW} steps. By repeating these experiments D times, one can get a series of data $\{f_{id}\}$ with $d = 1, 2, \dots, D$ and $i = 1, 2, \dots, N$. From the data we can calculate the average flux and the flux fluctuation of a node i as

$$\langle f_i(T_{RW}) \rangle = \frac{1}{D} \sum_{d=1}^D f_{id}(T_{RW}), \quad (2)$$

$$\sigma_i^2(T_{RW}) = \frac{1}{D} \sum_{d=1}^D f_{id}^2(T_{RW}) - \langle f_i(T_{RW}) \rangle^2. \quad (3)$$

From this model on the scale-free networks with the degree exponent $\gamma = 3$ the power-law relation $\sigma_i(T_{RW}) \simeq \langle f_i(T_{RW}) \rangle^\alpha$ or Eq. (1) was numerically shown. Furthermore the systems for small ΔN_{RW} were shown to belong to $\alpha \simeq 1/2$ class, while those for large ΔN_{RW} were shown to satisfy $\alpha = 1$. From these results it was explained why $\alpha = 1$ systems are governed by the external flux variations and the behavior for $\alpha = 1/2$ is from the internal intrinsic dynamics [2, 3].

In the multi-random-walk model, the flux configuration $\{f_{id}(T_{RW})\}$ exactly corresponds to the height configuration $\{h_{id}(T_{RW})\}$ in the following surface growth model. In the growth model each of N_{RW} random walkers drops a particle on the node i whenever it visits the node i and thus $f_{id}(T_{RW}) = h_{id}(T_{RW})$. Recently this kind of surface growth model in which N_{RW} is fixed (not-varied) in any experiment d is studied to know the effect of random-walk-like colored noises [7] on the dynamical scaling properties of the surface roughening [8]. In the study [7] the support is an ordinary one-dimensional lat-

*Electronic address: ykim@khu.ac.kr

tice and the dynamical scaling of the surface width W satisfies the conventional scaling as [8]

$$W = L^{\alpha_s} f(T_{RW}/L^z) = \begin{cases} T_{RW}^{\beta}, & T_{RW} \ll L^z \\ L^{\alpha_s}, & T_{RW} \gg L^z \end{cases} \quad (4)$$

with $z \equiv \alpha_s/\beta$.

In this paper we want to introduce another method to discriminate the systems governed by the internal dynamics from those derived by external input flux variations by comparison to corresponding surface growth dynamics. The main quantity for the comparison is the averaged $\sigma_i^2(T_{RW})$ over the node i , which is defined as

$$\bar{\sigma}^2(T_{RW}) \equiv \frac{1}{N} \sum_{i=1}^N \sigma_i^2(T_{RW}). \quad (5)$$

As we shall see, $\bar{\sigma}(T_{RW})$ for the multi-random-walker model shows the power-law dependence on the walk step T_{RW} as $\bar{\sigma}(T_{RW}) \simeq T_{RW}^{\beta}$. Furthermore, if the governing dynamics is not anomalous but the internal intrinsic dynamics, then $\bar{\sigma}(T_{RW})$ essentially has the same dynamical behavior as that of the mean-square surface fluctuation (width) $W(T_{RW})$ in one-random-walker surface growth model. Then T_{RW} is (acts as) the growth time and the exponent β is (or act as) the growth exponent in dynamical surface scaling [8]. If $W(T_{RW})$ in one-random-walker model dynamically behaves as $W(T_{RW}) \simeq T_{RW}^{\beta_i}$, then $\bar{\sigma}(T_{RW})$ for the systems governed by the internal intrinsic dynamics satisfies $\bar{\sigma}(T_{RW}) \simeq T_{RW}^{\beta}$ with $\beta = \beta_i$. If $\beta \neq \beta_i$, then governing dynamics should come from the external flux variations.

We first want to show the exact mathematical relation of $\bar{\sigma}$ to the surface fluctuation W . From Eq. (3),

$$\begin{aligned} \bar{\sigma}^2(T_{RW}) &= \frac{1}{N} \sum_{i=1}^N \sigma_i^2(T_{RW}) \\ &= \frac{1}{N} \sum_i \left[\frac{1}{D} \sum_{d=1}^D f_{id}^2 - \left(\frac{1}{D} \sum_{d=1}^D f_{id} \right)^2 \right]. \end{aligned} \quad (6)$$

Since $f_{id} = h_{id}$ in the corresponding surface growth model, the mean-square surface width W^2 is

$$\begin{aligned} W^2(T_{RW}) &= \frac{1}{D} \sum_{d=1}^D W_d^2(T_{RW}) \\ &= \frac{1}{D} \sum_d \left[\frac{1}{N} \sum_{i=1}^N f_{id}^2 - \left(\frac{1}{N} \sum_{i=1}^N f_{id} \right)^2 \right] \end{aligned} \quad (7)$$

Thus

$$\begin{aligned} \bar{\sigma}^2(T_{RW}) - W^2(T_{RW}) &= \frac{1}{D} \sum_d \left(\frac{1}{N} \sum_i f_{id} \right)^2 \\ &\quad - \frac{1}{N} \sum_i \left(\frac{1}{D} \sum_d f_{id} \right)^2. \end{aligned} \quad (8)$$

The difference between $\bar{\sigma}^2$ and W^2 for the given step T_{RW} comes from the different orders of taking the average over nodes (i.e., $1/N \sum_i f_{id}(T_{RW})$) and that over experiments (or trials) (i.e., $1/D \sum_d f_{id}(T_{RW})$).

In normal processes where any particular node does not receive extraordinarily smaller or larger flux than any other regions, then one can expect that the order of taking averages makes no anomalous effects and thus $1/N \sum_i f_{id}(T_{RW}) \simeq 1/D \sum_d f_{id}(T_{RW})$. Then the difference $\bar{\sigma}^2(T_{RW}) - W^2(T_{RW})$ becomes negligible or $\bar{\sigma}^2(T_{RW}) \simeq W^2(T_{RW})$. Therefore the possible origins which make the physically important difference comes from the breakdown of the spatial or temporal symmetry of the given system. Typical examples for such breakdowns are the local columnar defect [8, 10] and the temporally colored noises [8, 11]. When the input flux fluctuation becomes large or ΔN_{RW} is large in the multi-random-walker model, it is also possible that the input flux to some localized region of the support can become anomalously larger or smaller and then the difference $\bar{\sigma}^2(T_{RW}) - W^2(T_{RW})$ can have some crucial effects and remains for a certain length time.

To see the explained effects numerically, we analyze the numerical data from the simulations for the random-walker models. To see the universal features on homogeneous networks as well as inhomogeneous networks, we use regular lattices of one and two dimensions as well as random network(RN) [1] and scale-free network (SFN) with $\gamma = 3$ [1] as supports in the simulations. Used size or the number N of nodes (sites) of RN, SFN, and an one-dimensional lattice are 10^4 ($N = 10^4$). The used size of the two-dimensional square lattice is 100×100 . Each experiment (trial) is repeated over $D = 100$ times. Every walker in each experiment d takes 10^4 steps. Data for the flux $\{f_{id}(T_{RW})\}$ are obtained for the walk steps $T_{RW} = 1, 2, \dots, 10^4$.

The first simulation data to analyze are those for one-random-walker model in which only one random walker is initially dropped on a randomly chosen node (site) of the support in a given experiment. In this model the flux dynamics should be governed by the internal intrinsic dynamics and the difference $\bar{\sigma}^2(T_{RW}) - W^2(T_{RW})$ is expected to be negligible as explained in the previous paragraph. In Fig. 1 we have plotted the surface fluctuation $W(T_{RW})$ and the average flux variation $\bar{\sigma}(T_{RW})$ in one-random-walker model. On the four kinds of support (RN, SFN, one and two dimensional lattices) the dependence of $\bar{\sigma}(T_{RW})$ on T_{RW} (triangles in upper plots of Figs. 1(a)-(d)) is almost the same as that of $W(T_{RW})$ (straight lines in the same corresponding figures.). We can't find any significant difference $\bar{\sigma}^2(T_{RW}) - W^2(T_{RW})$. Furthermore the data for one random walker model satisfies the scaling ansatz $\bar{\sigma}(T_{RW}) = W(T_{RW}) = T_{RW}^{\beta_i}$ very well on both inhomogeneous networks (RN and SFN) and homogeneous networks (regular lattices). The estimated growth exponents β_i for one-random-walker model are $\beta_i = 0.53(2)$ for SFN, $\beta_i = 0.51(1)$ for RN, $\beta_i = 0.763(1)$ for one dimensional lattices and $\beta_i = 0.56(1)$ for two di-




FIG. 1: Triangles ($\blacktriangle$) and straight lines in the upper plots of figures (a), (b), (c), and (d) represent the data for the average flux fluctuation $\bar{\sigma}(T_{RW})$ over nodes and the surface fluctuation $W(T_{RW})$ in the one-random-walker model, respectively. Other symbols in the upper plots represent $\bar{\sigma}(T_{RW})$ for $\Delta N_{RW} = 0, 20, 40, 80, 100, 200, 800, 1000, 4000, 10000$ from bottom to top in the multi-random walker model. The lower plots show the values of exponent β for various ΔN_{RW} obtained by fitting the data in upper plots to the relation $\bar{\sigma}(T_{RW}) \simeq T_{RW}^\beta$. The dotted lines in the lower plots denote the values of β_i obtained from the relation $W(T_{RW}) \simeq \bar{\sigma}(T_{RW}) \sim T_{RW}^{\beta_i}$ in one-random-walker model. (a) on Scale-free network with $\gamma = 3$. (b) on Random network. (c) on one dimensional lattice. (d) on two dimensional lattice.

mensional lattices. On the inhomogeneous networks such as SFN and RN, β_i is very close to $\beta = 1/2$, which is the value of β of random ballistic depositions on any support [8]. In contrast on one and two dimensional lattices β_i is quite different from $1/2$.

In the simulations for multi-random-walker models, the number N_{RW} of walkers in an experiment d is selected randomly among the uniform distribution

$[\langle N_{RW} \rangle - \Delta N_{RW}, \langle N_{RW} \rangle + \Delta N_{RW}]$. Then the N_{RW} walkers are initially randomly distributed over the nodes (sites) of the support. In the simulations $\langle N_{RW} \rangle = 10^4$ is always imposed and ΔN_{RW} is varied as $\Delta N_{RW} = 0, 20, 40, 80, 100, 200, 800, 10^3, 4 \times 10^3, 10^4$. Other conditions are exactly the same as those in the simulations for one-random-walker model. The results for the dependence of $\bar{\sigma}$ on T_{RW} in multi-random-walker model for various ΔN_{RW} are also shown in upper plots of Figs. 1(a)-(d). From the data and the relation $\bar{\sigma} \simeq T_{RW}^\beta$ the obtained values of β for ΔN_{RW} 's are displayed in the lower plots of Figs. 1(a)-(d), where the dotted line denotes the values of β_i in one-random-walker model on the corresponding support. As can be seen from the lower plots, β value for small ΔN_{RW} or $\Delta N_{RW} \leq 200$ is nearly equal to β_i (i.e., $\beta = \beta_i$) on both homogenous and inhomogeneous networks. On the other hand β for large ΔN_{RW} or $\Delta N_{RW} \geq 4 \times 10^3$ approaches to $\beta = 1$ (or $\beta \geq 0.9$). β for crossover values of ΔN_{RW} (or $\Delta N_{RW} = 800$ or 1000) has crossover values between β_i and 1. The results for the multi-random walker models are as follows. For small ΔN_{RW} or in the systems where the internal intrinsic dynamics is dominant, $\bar{\sigma}(T_{RW})$ follows the behavior of $W(T_{RW}) \simeq T_{RW}^{\beta_i}$ in one-random-walker model quite well. For large ΔN_{RW} or in the systems where the external flux fluctuation is dominant for dynamical behaviors, $\bar{\sigma}(T_{RW})$ follows the behavior $\bar{\sigma}(T_{RW}) \simeq T_{RW}^\beta$ with $\beta = 1$.

In summary, we have shown that $\bar{\sigma}$ shows the power-law dependence on the walk step T_{RW} as $\bar{\sigma} \simeq T_{RW}^\beta$ in multi-random-walker model for various ΔN_{RW} . To discriminate the systems governed by the internal intrinsic dynamics from those governed by the external flux variations, the dynamical behavior of one-random-walker model is suggested as the criterion. If the dependence of $\bar{\sigma}$ in multi-random-walker model on T_{RW} follows the dynamical behavior of the surface width in one-random-walker model as $W(T_{RW}) \simeq T_{RW}^{\beta_i}$ or $\beta = \beta_i$, then the model is governed by the internal dynamics. This behavior occurs when ΔN_{RW} is small. When $\beta \neq \beta_i$ and $\beta = 1$, then the model is governed by the external flux variations. The $\beta = 1$ behavior occurs when ΔN_{RW} is large. We also show that this criterion holds for the model not only on the inhomogeneous networks but also on homogeneous networks or regular lattices. Furthermore β_i for one and two dimensional lattices is greater than $1/2$. Especially in one dimension the random walker has the temporal correlation due to the reentrant property and the visitation probability to a certain site has some power-law correlation in time [7, 9]. In two-dimension the reentrant property is marginal [9]. This reentrant property of random walker explains $\beta_i > 1/2$ in both one and two dimensional lattices. In contrast for SFN and RN whose effective dimensionality is ∞ [1], the random walkers are transparent [9] and the one-random-walker model behave as random ballistic deposition and satisfies $\beta = 1/2$ [8].

Final comments are on the nonuniversal values of exponent α for Eq. (1) in regular lattices or homogeneous net-




FIG. 2: Dependence of the exponent α on ΔN_{RW} of multi-random-walker model in one (1d) and two (2d) dimensional lattices. The data is obtained by fitting $\sigma_i(T_{RW})$ versus $\langle f_i(T_{RW}) \rangle$ to the relation (1) or $\sigma_i(T_{RW}) \simeq \langle f_i(T_{RW}) \rangle^\alpha$. The data for $\sigma_i(T_{RW})$ and $\langle f_i(T_{RW}) \rangle$ in one and two dimensions are obtained from the same simulation conditions as those in Figs. 1(c) and (d) with a fixed value of T_{RW} as $T_{RW} = 100$.

works. Recently a study [4] suggested a model in which nonuniversal α values (i.e., $\alpha \neq 1$ or $1/2$) for the relation $\sigma_i(T_{RW}) \simeq \langle f_i(T_{RW}) \rangle^\alpha$ with the fixed T_{RW} . By using the multi-random-walker model with the quenched impact variable $V(i)$ for each node i , which is dependent

on the degree $k(i)$, the study [4] showed nonuniversal α values (i.e. $1/2 < \alpha < 1$) on scale-free networks. Nonuniversal α values are also possible on the homogeneous networks or regular lattices without the impact variable $V(i)$. By using the relation $\sigma_i(T_{RW}) \simeq \langle f_i(T_{RW}) \rangle^\alpha$ with the fixed T_{RW} on one and two dimensional lattices, we obtain α values for various ΔN_{RW} . The results are shown in Fig. 2. The simulation conditions except T_{RW} for Fig. 2 are exactly the same as those for Figs. 1(c) and (d). T_{RW} for each experiment (day) is fixed as $T_{RW} = 100$. For $\Delta N_{RW} < 500$ where the internal intrinsic dynamics are dominant, α values are nearly constant. The values of the exponent α for small ΔN_{RW} are $\alpha = 0.77(1)$ in one dimension and $\alpha = 0.60(2)$ in two dimension, respectively. The behavior $\alpha > 1/2$ for internal intrinsic dynamics should come from the reentrant properties of random walks. As explained before the reentrant property makes the power-law correlation in time for the probability to visit a certain site again. This kind of temporal correlation effects makes the probability distribution for $\langle f_i \rangle$ deviate from the Gaussian and thus gives nonuniversal α values for small ΔN_{RW} . For large ΔN_{RW} α approaches 1 even for homogeneous networks as can be seen from data for $\Delta N_{RW} = 4000, 10000$ in Fig. 2.

This work was supported by Grant No. R01-2004-000-10148-0 from the Basic Research Program of KOSEF. We thank Dr. Soon-Hyung Yook for valuable suggestions. One of authors (YK) also acknowledges the hospitality of Korea Institute of Advanced Study during the visit.

-
- [1] R. Albert and A.-L. Barabási, Rev. Mod. Phys. **74**, 47 (2002); S.N. Dorogovtsev and J.F.F. Mendes, Adv. Phys. **51**, 1079 (2002).
 - [2] M. Argollo de Menezes and A. -L. Barabási, Phys. Rev. Lett. **92**, 028701 (2004).
 - [3] M. Argollo de Menezes and A. -L. Barabási, Phys. Rev. Lett. **93**, 068701 (2004).
 - [4] Z. Eisler and J. Kertézes, Phys. Rev. E **71**, 057104 (2005).
 - [5] Z. Eisler and J. Kertézes, S. -H. Yook, and A. -L. Barabási, arXiv:cond-mat/0408409 (2004).
 - [6] Z. Eisler and J. Kertézes, arXiv:cond-mat/0503139 (2005).
 - [7] H. S. Song and J. M. Kim, Jour. of the Korean Phys. Soc. **44**, No. 3, 543 (2004).
 - [8] *Dynamics of Fractal Surfaces*, edited by F. Family and T. Vicsek (World Scientific, Singapore, 1991); A.-L. Barabási and H. E. Stanley, *Fractal Concepts in Surface Growth* (Cambridge University Press, Cambridge, 1995); J. Krug, Adv. Phys. **46**, 139 (1997).
 - [9] S. Redner, *A GUIDE TO FIRST PASSAGE PROCESSES* (Cambridge University Press, Cambridge, 2001).
 - [10] T. Halpin-Healy and Y.-C. Zhang, Phys. Rep. **254**, 215 (1995).
 - [11] E. Medina, T. Hwa, M. Kardar and Y.-C. Zhang, Phys. Rev. A **39**, 3053 (1989).